

## Age, sex, religion and caste specific goitre prevalence among school children in Berhampore sadar sub-division of Murshidabad District, West Bengal

Smritiratan Tripathy<sup>1\*</sup>, Gopal Chandra Halder<sup>2</sup> and Arijit Debnath<sup>3</sup>

<sup>1</sup>Department of Physiology, Berhampore Girls' College, P.O. Berhampore, Dist. Murshidabad West Bengal, India, <sup>2</sup>Hasanpur Rejeswari Vidyapith, Hasanpur, Murshidabad, West Bengal, India and <sup>3</sup>Krishnabati High School, Simuliadanga, Burdwan, West Bengal, India

**Abstract:** *Objectives:* The objective of the present study is to find out the prevalence of endemic goitre among school children of Berhampore sadar sub-division of Murshidabad district. Attempt has also been made to find out the interaction of host factors, viz, age, sex, caste, religion in influencing goitre prevalence among school children during post-salt iodization phase. *Background:* In post salt iodization phase, endemic goitre was found prevalent all over the India including West Bengal. Prevalence of goitre has been reported in many districts except Murshidabad. Present study is thus undertaken in Murshidabad district. *Method:* A total of 1566 school children (6-12 yr) were clinically examined for goitre from five C.D.Blocks of the studied region. Age, sex, religion and caste specific goitre prevalence was reported. Body height and body weight is also recorded to observe the growth pattern of the studied school children. *Result:* Overall result showed that mild degree of goitre endemicity (17.4%) was found. Girls (18.9%) were more affected than boys (15.7%). Goitre was most prevalent in Hindus (19.6%) than Muslim (16.0%) communities. Among the Hindus, general caste (20.4%) students suffered more than scheduled caste and scheduled tribe (18.6%). *Conclusion:* Present study indicates that goitre is still persist in the region. Result showed that girls were more affected than boys. It may be due to increase physiological demand in females at puberty. Result also showed that goitre was most prevalent in Hindus than Muslim communities and among the Hindus, general caste students suffered more than scheduled caste & scheduled tribe. It might be due to their different environmental adaptability and different dietary habit. Age specific data showed that in initial stage (6 yr) growth pattern is comparably better in boys than girls. The growth rate is high among the girls than boys at the age of 9 yr to 12 yr. It may be due to the high demand of steroid hormones among the girls before menarche.

**Keywords:** Goitre, School children, Murshidabad

### Introduction

Iodine deficiency disorders are still a major public health problem in India. The major consequences of iodine deficiency are goitre (enlargement of thyroid gland than normal), mental defect, deaf mutism, stillbirth and miscarriage, weakness and paralysis of muscles as well as lesser degree of physical and mental function [1]. Effects on brain function occur at all stages of life, from foetal damage or hypothyroidism in the neonate, child or adult. Considerable progress has been made in the implementation of the universal salt iodization programme in the countries affected by IDD. IDD is now not restricted in the classical hilly sub-Himalayan goitre endemic belt in india. It has also been reported from sub-Himalayan flat lands (tarai), plains (especially those subjected to annual

flooding), riverine areas, deltas and even in coastal regions [2-3]. The northern part of the state of West Bengal is hilly and located in the classical iodine deficient goitre endemic belt in India [4], while its major southern region is in the Gangetic basin where the land is plain, fertile and thickly populated. Prevalence of goitre has been reported in many districts like South 24-Parganas, North 24-Parganas, Howrah, Malda, Purulia, Dakhin Dinajpur and Birbhum in West Bengal [5-12]. The systemic studies about the prevalence of goitre among school children of the Gangetic regions of Murshidabad district are not available. Present study is thus undertaken to evaluate the present state of goitre prevalence among the school children of Berhampore sadar Sub-division in Murshidabad district as per

WHO/UNICEF/ICCIDD recommendation. In the present study attempt has also been made to find out the interaction of host factors, viz, age, sex, caste, religion in influencing goitre prevalence among school children during post-salt iodization phase.

**Material and Methods**

Murshidabad district is a district of West Bengal in Eastern India. It is one of the 18 district of West Bengal. There are 26 C.D.Blocks under 5 sub-division in Murshidabad district. The study region Berhampore sadar subdivision is one of them consisting with 5 C.D.Blocks. Murshidabad district is situated on the left bank of the river Ganges. Total area of the district is 5341 sq.km. This district is well known for his historical importance. The plight of whole Bengal changed with the plight of Murshidabad. But at present despite there has been much more progress of West Bengal in every way but there is no significance change of Murshidabad district. Poverty and illiteracy have not been eradicated to the possible extent. Most of the people are engaged in agricultural activities. To get proper representation five areas were selected from five C.D. Blocks taking one from each by random purposive sampling method [13]. In each selected areas one primary school and nearest adjoining secondary school were selected at random [6] where the students of the age group 6-12 yr. of both sexes were available as recommended by WHO/UNICEF/ICCIDD [14]. The entire study was conducted in between February 2011 and November 2011. A total of 1566 students were clinically examined for goitre. Goitre grading was done according to the criteria recommended by joint WHO/UNICEF/ICCIDD (1994) (grade 0: no

goitre; grade 1: thyroid palpable but not visible; and grade 2: thyroid visible with the neck in normal position) [15]. The age of the students was recorded from the school register and was rounded off to the nearest whole number.

Anthropometric data mainly height and weight of a national population or a particular segment represent its health status in relation to heredity and environment. In iodine deficient region physical growth is retarded. To evaluate the physical growth pattern height and weight pattern is also recorded. The age of the subjects was determined from their dates of birth recorded in the school register. The age was rounded off to the nearest whole number (e.g., age of the students ranging from 13.50 to 14.49 yr was expressed as 14+). Standing body height was measured by anthropometric rod and weight was taken by standard weighing machine. All measurements were taken according to the method described by Damon et al [16].

**Results and Discussion**

Prevalence of endemic goitre in school children is the most widely accepted marker to evaluate the severity of IDD in the region [15]. According to WHO/UNICEF/ICCIDD recommended criteria, a prevalence rate of 5.0 – 19.9 % is considered mild; 20.0 – 29.9 % as moderate and above 30 % is considered as a severe public health problem [15]. The overall goitre prevalence of 17.4 % indicates that IDD is a mild degree of endemicity as a public health problem in the studied region (Table 1).

**Table-1: Goitre prevalence in different study areas of Berhampore sadar subdivision in Murshidabad district**

Sl No.	Study areas (C.D. Blocks)	Total number of children examined	Number of children with goitre			Severity as public health problem
			Grade-1	Grade-2	Total (1+2)	
1	Berhampore	263	53 (20.1)	12 (4.6)	65 (24.7)	Moderate
2	Beldanga - I	299	45 (15.0)	04 (1.3)	49(16.4)	Mild
3	Beldanga - II	353	60(17.0)	05 (1.4)	65 (18.4)	Mild
4	Hariharpara	383	55 (14.4)	04(1.0)	59 (15.4)	Mild
5	Nawda	268	30(11.2)	04 (1.5)	34 (12.7)	Mild
<b>Total</b>		<b>1566</b>	<b>243(15.5)</b>	<b>29 (1.8)</b>	<b>272 (17.4)</b>	<b>Mild</b>

Severity of public health problem: 5.0-19.9% mild; 20.0-29.9% moderate; >30% Severe (Parentheses indicate percentage)

**Table-2: Prevalence of goitre in boys, girls, and overall studied population of Berhampore sadar sub-division of Murshidabad district.**

Age (years)	BOYS				GIRLS				TOTAL			
	No. of Children examined	Grades of Goitre			No. of children examined	Grades of Goitre			No. of children examined	Grades of Goitre		
		1	2	Total		1	2	Total		1	2	Total
6	150	13	2	15 (10.0)	180	30	-	30 (16.7)	330	43	2	45 (13.6)
7	134	29	2	31 (23.1)	156	27	-	27 (17.3)	290	56	2	58 (20.0)
8	144	19	1	20 (13.9)	140	20	2	22 (15.7)	284	39	3	42 (14.8)
9	123	23	3	26 (21.1)	98	10	2	12 (12.2)	221	33	5	38 (17.2)
10	106	17	2	19 (17.9)	109	16	8	24 (22.0)	215	33	10	43 (20.0)
11	69	09	-	9 (13.0)	108	30	7	37 (34.2)	177	39	7	46 (26.0)
12	37	-	-	-	12	-	-	-	49	-	-	-
<b>Total</b>	763	110 (14.4)	10 (1.3)	120 (15.7)	803	133 (16.6)	19 (2.4)	152 (18.9)	1566	243	29	272 (17.4)

(Parentheses indicate percentage)

**Table-3: Age specific religion wise goitre prevalence among school children of Berhampore sadar sub-division of Murshidabad district**

Age (years)	HINDUS				MUSLIM			
	No. of Children examined	Grades of Goitre			No. of children examined	Grades of Goitre		
		1	2	Total		1	2	Total
6	142	24	2	26 (18.3)	188	19	-	19 (10.1)
7	92	14	-	14 (15.2)	198	42	2	44 (22.2)
8	112	19	2	21 (18.9)	172	20	1	21 (12.2)
9	88	19	2	21 (23.9)	133	14	3	17 (12.8)
10	73	8	3	11 (15.0)	142	25	7	32 (22.5)
11	68	18	7	25 (36.8)	109	21	-	21 (19.3)
12	28	-	-	-	21	-	-	-
<b>Total</b>	603	102 (16.9)	16 (2.6)	118 (19.6)	963	141 (14.6)	13 (1.3)	154 (16.0)

Parentheses indicate percentage

Though most of the goitre was found palpable (15.5%) but the prevalence of visible goitre (1.8%) among the children of 6-12 yr. also exist. The multifactorial nature and complex interactions of host factors (age, sex) with region-specific environmental conditions in the pathogenesis of endemic goitre constitute a major challenge in the understanding and control of the problem of goitrogenic substances in endemic areas [17]. Age specific sex wise goitre prevalence in overall studied students is given in Table 2. A gradual increase in goitre rate is found from the age of 6 yr till the age of 11 yr except a short decline in the age of 8 and 9 yr. Age wise overall goitre rates in boys and girls were in the ranges of 10% - 23% and 12.2% - 34.2% excluding the age group of 12 yr as the number of

sample is poor in this group. The overall result showed that girls (18.9%) were more affected in goitre than boys (15.7%). The prevalence of grade 2 goitre is also found higher in girls (2.4%) than boys (1.3%). The higher prevalence in females has been generally attributed to increase physiological demands at puberty [18]. Earlier works showed that thyroid disorders (endemic goitre) are common in women than men especially in iodine deficient region [19]. This is because estrogens appear to enhance the response to TRH (thyrotropin releasing hormone) possibly by increasing the number of TRH receptors in the thyrotrophic cells in pituitary [20].

**Table-4: Age specific caste wise goitre prevalence among Hindu school children of Berhampore sadar sub-division of Murshidabad district**

Age (years)	GENERAL				SC/ST			
	No. of Children examined	Grades of Goitre			No. of children examined	Grades of Goitre		
		1	2	Total		1	2	Total
6	75	19	-	19 (25.3)	67	5	2	7 (10.4)
7	53	6	-	6 (11.2)	39	8	-	8 (20.5)
8	37	6	-	6 (16.2)	75	13	2	15 (20.0)
9	41	5	-	5 (12.2)	47	14	2	16 (34.0)
10	45	6	2	8 (17.8)	28	2	1	3 (10.7)
11	38	13	6	19 (50.0)	30	5	1	6 (20.0)
12	19	-	-	-	9	-	-	-
Total	308	55 (17.8)	8 (2.6)	63 (20.4)	295	47 (15.9)	8 (2.0)	55 (18.6)

Parentheses indicate percentage

Overall goiter prevalence in respect of religion is shown in Table 3. It showed that goitre was most prevalent in Hindus (19.6%) than Muslim (16.0%) communities. Among the Hindus, general caste (20.4%) students suffered more than Scheduled caste & Scheduled tribe (18.6%) (Table 4). The observed result is almost

consistent with the earlier observation of Chandra and Ray (2001) [21]. The people of all caste and religion were exposed under uniform system of iodine supply and consume foods namely beans, turnip, sweet potato, cabbage, cauliflower, radish, mustard, etc containing naturally occurring goitrogens

available in the region. But these variations in goitre prevalence among different castes might be due to their different environmental adaptability and dietary habit, as has been observed in a similar study conducted in Africa by Biassoni et al, 1998 [22], where it was found that minute difference in iodine deficiency between two populations induces to tally different pattern of goitre and thyroid function probably for the

differences in diet. Body weight and body height of the studied school children is shown in Table 5. Age specific data showed that in initial stage (6 yr) growth pattern is comparably better in Boys than girls. The growth rate is high among the girls than boys at the age of 9 yr to 12 yrs. It may be due to the high demand of steroid hormones among the girls before menarche.

**Table-5: Age and sex specific body weight and body height of school children of Berhampore sadar sub-division of Murshidabad district (Data are only mean)**

Age (years)	BOYS			GIRLS		
	No. of Children Examined	Weight (Kg)	Height (c.m.)	No. of Children Examined	Weight (Kg)	Height (c.m.)
6	150	15.79	107.24	180	14.96	106.89
7	134	17.93	113.67	156	16.87	113.63
8	144	19.27	118.46	140	19.06	118.64
9	123	20.38	121.8	98	21.02	122.56
10	106	22.5	125.79	109	21.71	124.46
11	69	23	128.54	108	23.62	129.02
12	37	28	132.61	12	28.25	134.75

The present result indicates that goitre is still persist in this region. To prevent and control this public health problem, it is essential to observe the present state of iodine nutritional status in the studied region.

**Acknowledgements**

The financial assistance by the University Grants Commission (PSW-080/10-11) is gratefully acknowledged. The authors acknowledge the cooperation received from the staff and students of the schools studied.

**References**

- Hetzel BS An overview of the prevention and control of iodine deficiency disorders. In: Hetzel BS, Dunn JT, Stanbury JB editors. The prevention and control of iodine deficiency disorders. Elsevier, Amsterdam, 1987; 7-31.
- Kochupillai N. Neonatal hypothyroidism in India. *Mount Sinai Journal of Medicine*, 1992; 59(2):111-5.
- Pandav CS, Malik A, Anand K, Pandav S and Karmarkar MG. Prevalence of iodine deficiency disorders among school children of the national capital territory of Delhi. *ICCIDD, Regional Office of South Asia and Pacific, New Delhi*; 1996.
- WHO/SEARO. Iodine deficiency disorder in Southeast Asia. SEARO Regional Health, New Delhi, Paper No.10; 1985.
- Chandra AK, Tripathy S, Lahari D, Mukhopadhyay S. Iodine nutritional status of the school children in a rural area of Howrah District in the Gangetic West Bengal. *Indian J Physiol & Pharmacol* 2004; 48(2): 219-224
- Chandra AK, Tripathy S, Ghosh D, Debnath A, Mukhopadhyay S. Iodine nutritional status & prevalence of goitre in Sundarban delta of South 24-Parganas, West Bengal, *Indian J Med Res* 2005;122 (5): 419-424.
- Chandra AK, Tripathy S, Ghosh D, Debnath A, Mukhopadhyay S. Goitre prevalence and state of iodine nutrition in Sundarban delta of North 24-Parganas in West Bengal, *Asia Pac J Clin Nutr* 2006;15 (3):357-361.
- Chandra AK, Debnath A, Tripathy S. Iodine nutritional status among school children in selected areas of Howrah district in West Bengal, India. *J Trop Pediatr*. 2008; 54(1): 54-57.
- Biswas AB, Chakraborty I, Das DK, Biswas S, Nandy S, Mitra J. Iodine deficiency disorders among school children of Malda, West Bengal, India. *J Health Popul Nutr* 2002; 20(2): 180-183.
- Biswas AB, Chakraborty I., Das DK, Roy RN, Mukhopadhyay S, Chatterjee S. Iodine deficiency disorders among school children of Birbhum, West Bengal. *Current Science* 2004; 87(1): 78-80.
- Das DK, Chakraborty I., Biswas AB, Sarkar GN, Srivastava P, Sen S. Iodine deficiency disorders among school children of Dakshin Dinajpur

- District, West Bengal. *Indian J Public Health* 2005; 49(2): 68-72.
12. Biswas AB, Chakraborty I, Das DK, Roy RN, Ray S, Kunti SK. Assessment of iodine deficiency disorders in Purulia district, West Bengal, India. *J Trop Pediatr* 2006; 52(4): 288-292.
  13. Cochran WG. *Sampling technique*, 3<sup>rd</sup> ed. Wiley Eastern Limited, Calcutta; 1977.
  14. WHO/ UNICEF/ ICCIDD. Assessment of the Iodine Deficiency Disorders and Monitoring their Eliminations. Geneva: WHO Publ, WHO/NHD/01.1.1-107; 2001.
  15. WHO/UNICEF/ICCIDD. Indicators for assessing Iodine Deficiency Disorders and their control through salt iodization. WHO/NUT/94.6; 1994.
  16. Damon A, Stoudt HW and McFarland RA. The human body in equipment design (Harvard University Press, Cambridge), 1996; 62.
  17. Gaitan E and Dunn JT. Epidemiology of iodine deficiency. *Trends Endocrinol Metab*, 1992; 3:170-175.
  18. Dudani TG and Natu MN. Epidemiology of endemic goitre in Ghodegaon. *Indian J Med Res.* 1978; 68: 980-989.
  19. Glinoeer D and Leome M. Goitre and pregnancy: a new insight into an old problem. *Thyroid* 1996; 2: 65-69.
  20. Gershengorn MC, Marcus-Samuels BE and Geras F. Estrogens increase the number of thyrotropin releasing hormone receptors on mammotropic cells in culture. *Endocrinology*. 1979; 105: 171.
  21. Chandra AK and Ray I. Influence of age, sex and caste on goiter prevalence of the people in Tripura, North East India. *J Human Ecology*; 2001; 2(4): 313-317.
  22. Biassoni P, Ravera G, Bertocchi J, Schenone F and Bourdoux P. Influence of dietary habits on thyroid status of a nomadic people, the Bororo shepherd roaming a certain African regions affected by severe iodine deficiency. *Eur J Endocrinol*; 1998; 138: 681-685.

\*All correspondences to: Dr. Smritiratan Tripathy, Assistant Professor, Department of Physiology, Berhampore Girls' College, P.O. Berhampore, Dist. Murshidabad, West Bengal-742101 India. E-mail: smritiratan\_tripathy@yahoo.com