

## Anthropometric, physical and cardiorespiratory fitness of 10-16 years children

Indranil Manna<sup>1\*</sup>, Swadesh Ranjan Pan<sup>2</sup> and Mohua Chowdhury<sup>3</sup>

<sup>1</sup>Department of Physiology, Midnapore College, Raja Bazar Main Road, Midnapore, West Bengal, India, <sup>2</sup>Department of Physical Education, Jhargram Raj College, Jhargram, Paschim Medinipur, West Bengal, India and <sup>3</sup>Department of Community Medicine, Bankura Sammilani Medical College, P.O. Kenduadihi, Bankura, West Bengal, India

**Abstract:** *Objectives:* The present study was undertaken to investigate the anthropometric, physical and cardiorespiratory fitness of 10-16 yrs children. *Background:* Talent identification in sports is importance because they represent the achievement level of a particular group in future. There are very limited studies available in Indian context on talent identification in sports. *Method:* A total of 150 male children of 10-16 yrs age volunteered for this study; were divided equally into 3 groups (i) Prepubertal (age-11.0 ± 0.8yrs, n=50); (ii) Pubertal (age 13.5 ± 0.5 yrs, n=50); (iii) Postpubertal (age 15.5 ± 0.5 yrs, n=50). Selected anthropometric, physical and cardiorespiratory fitness variables were measured for each group. *Result:* A significantly (P<0.05) greater height, body mass, BSA, LBM, mid upper arm circumference, hip and trunk flexibility, grip strengths, abdominal strength, elastic leg strength, maximum speed, peak power, VO<sub>2max</sub>, peak expiratory flow rate and blood pressure were observed in Postpubertal children when compared to Prepubertal and Pubertal children. However, a significantly (P<0.05) lower percent body fat, reaction time, maximal heart rate and recovery heart rates were noted in Postpubertal children when compared to Prepubertal and Pubertal children. The waist- hip ratio of pubertal children was noted significantly higher (P<0.05) when compared to prepubertal and postpubertal children. No significant change was reported in BMI and resting heart rate among the groups. *Conclusion:* Identification of children at early stage of their growth and development may produce elite athletes in the future. Talent identification also can be used as a counseling technique that helps to discover and explore areas of talent for particular athletes.

**Keywords:** body composition, VO<sub>2max</sub>, power, strength, heart rate.

### Introduction

Sports talent may be identified from the school children 10-16 yrs age group when they show interest in different sports. Identification and selection of talented children for sports are not straightforward operations [1]. In developed countries like USA and Australia identification of sports talent is performed scientifically. However, lack of scientific knowledge and infrastructural facilities for identification of sports talent, result into poor performance of Indian athletes at international arena. Anthropometric, physical and cardiorespiratory fitness profiles contribute to selection procedures in different sports events [2]. Besides success in track and field discipline is based on the synthesis of anthropometric characteristics and motor abilities as well as optimal technique [3]. But overall characteristics are also influenced by genetic inheritance, morphology, personal interest and habitual

activity. Cardiorespiratory fitness variables such as maximal aerobic capacity, heart rate, blood pressure and pulmonary functions reflect the overall capacity of the cardiovascular and respiratory systems and the ability to carry out prolonged exercise [4]. Hence, Cardiorespiratory fitness has been considered as a direct measure of the physiological status of the individual [4-5].

The gold standard for the measurement of cardiorespiratory fitness is the maximal oxygen uptake (VO<sub>2max</sub>). The level of cardiorespiratory fitness is highly associated with the performance of other health-related fitness parameters such as strength and power output in young people and in adults [6]. To identify athletic potentiality, norms of the anthropometric, physical and cardiorespiratory fitness profiles have an

importance because they represent the achievement level of a particular group to which observed score can be compared [7]. Various factors like socio-economic condition, diet, physical activity may reflect on these measurements. Thus there is a wide range of normalcy and the need to develop local norms has been emphasized. Several studies have been carried out on the physical [6] and cardiorespiratory [5] fitness status of the children of school-age populations. In India, limited studies on the anthropometric, physical, [8-10] and cardiorespiratory fitness [9-10] of children have been reported. In view of the above, a study was undertaken to investigate the anthropometric, physical and cardiorespiratory fitness of 10-16 years age group children in order to identify potentiality and sports talent in them.

### Material and Methods

*Subjects:* A total of 150 male children of 10-16 yrs age volunteered for this study. The children were selected after proper medical checkups from West Midnapore districts of West Bengal, India. The subjects were equally divided into 3 groups (i) Prepubertal (age-11.0  $\pm$  0.8yrs, n=50); (ii) Pubertal (age 13.5  $\pm$  0.5 yrs, n=0); (iii) Postpubertal (age 15.5  $\pm$  0.5 yrs, n=50). The subjects were informed about the possible complications of the study and gave their consent. Parental consent was also taken from the participants of this study. The institutional review board and ethical board approval was also obtained for the present study.

*Measurement of Anthropometric Variables:* Height and body mass were measured using standard methodology [11]. Body mass index (BMI) and Body surface area (BSA) were derived from the height and body mass using standard equations [11]. Measurements of hips and waist of the subject was taken by a steel tape using standard procedure, and the waist- hip ratio (WHR) was determined by standard equation [11]. Mid upper arm circumference (MUAC) of the subject was taken by a steel tape using standard procedure [11]. A skin fold calliper (Mitutoyo, Japan) was used to assess the body fat percentage, from biceps, triceps, sub scapular and suprailiac sites. Body density was calculated according to the formulae of Durnin and Womersley [12]. Body fat was derived using the standard equation of Siri [13]. Subsequently, lean

body mass (LBM) was derived by subtracting fat mass from total body mass using the standard equation [11].

*Assessment of Physical Fitness:* Reaction time of the subject was assessed by ruler drop test using standard procedure [11]. Modified sit and reach test (MSRT) was applied using standard procedure in order to assess subject's hip and trunk flexibility [11]. Sit ups test (SUT) was performed using standard procedure to monitor the development of the subject's abdominal strength [11]. Standing long jump test (SLJT) was performed to monitor the development of the subject's elastic leg strength [11]. A grip strength dynamometer (T.K.K.5001 Grip-A, Japan) was used to record the strength of grip muscles of both hands following a standard methodology [11]. To monitor the development of the subject's ability to effectively and efficiently build up acceleration from a standing start to maximum speed, 30 meter acceleration test (30MAT) was performed using standard procedure [11]. Margaria Kalamen Power Test was used to monitor subject's peak power using the standard procedure [14].

*Assessment of Cardiorespiratory fitness:* Subject was asked to take rest for 15 min and the heart rate and blood pressure were recorded. Maximal heart rate (HRmax) and recovery heart rates were recorded following a maximal exhaustive exercise. Maximal aerobic capacity ( $VO_{2max}$ ) was measured indirectly using Queen's College step test following standard procedure [15]. Peak expiratory flow rate (PEFR) was recorded using a vitalograph (Montari, India) following standard procedure [16].

*Statistical Analysis:* All the values of anthropometric, physical and cardiorespiratory fitness variables were expressed as mean and standard deviation (SD). Analysis of Variance (ANOVA) followed by multiple comparison tests was performed to find out the significant difference in selected anthropometric, physical and cardiorespiratory fitness variables among the groups. In each case the significant level will be chosen at 0.05 levels. Accordingly, a statistical software package (SPSS) was used.

**Results**

Anthropometric parameters showed variations among the Prepubertal, Pubertal and Postpubertal children. A significantly (P<0.05) higher height, body mass, body surface area (BSA), lean body mass (LBM) and mid upper arm circumference (MUAC) were observed in Postpubertal children when compared to Prepubertal and Pubertal children. On the other hand, a significantly

(P<0.05) lower percent body fat was noted in Postpubertal children when compared to Prepubertal and Pubertal children. The waist-hip ratio (WHR) of pubertal children was noted significantly higher (P<0.05) when compared to prepubertal and postpubertal children. However, no significant change was reported in body mass index (BMI) and total body fat among the groups (Table 1).

**Table-1: Anthropometric variable of Prepubertal, Pubertal and Postpubertal children**

Variables	Prepubertal	Pubertal	Postpubertal
Age (yr)	11.0 ± 0.8	13.5 ± 0.5*	15.5 ± 0.5*#
Height (cm)	140.2 ± 4.1	152.5 ± 4.2*	161.7 ± 4.3*#
Body mass (kg)	34.8 ± 5.2	41.9 ± 4.2*	46.5 ± 4.5*#
BMI	17.3 ± 1.5	18.0 ± 1.5 <sup>NS</sup>	17.8 ± 1.2 <sup>NS</sup>
BSA	1.2 ± 0.04	1.3 ± 0.05*	1.5 ± 0.04*#
Body fat (%)	18.6 ± 2.4	17.3 ± 2.1*	15.0 ± 2.7*#
Body Fat (kg)	6.8 ± 1.3	7.3 ± 1.6 <sup>NS</sup>	7.1 ± 1.5 <sup>NS</sup>
LBM (kg)	28.0 ± 4.3	34.6 ± 3.6*	39.4 ± 3.9*#
WHR	0.8 ± 0.02	0.9 ± 0.02*	0.8 ± 0.02#
MUAC (cm)	17.5 ± 2.9	20.8 ± 2.7*	23.2 ± 2.5*#

All the values were expressed as mean and standard deviation (SD), n=50; ANOVA followed by multiple comparison tests; \*P<0.05 when compare to prepubertal age group, #P<0.05, when compare to pubertal age group, NS= not significant; BMI= body mass index, BSA= body surface area, LBM= lean body mass, WHR= waist- hip ratio, MUAC= mid upper arm circumference.

Physical fitness variables showed remarkable differences among the Prepubertal, Pubertal and Postpubertal children. A significantly (P<0.05) higher hip and trunk flexibility as measured by modified sit and reach test (MSRT) score, abdominal strength as measured by sit ups test (SUT) score, elastic leg strength as measured by standing long jump test (SLJT) score, grip strengths of both hands, maximum speed as

measured by lower 30 meter acceleration test (30MAT) score, and peak power output were observed in Postpubertal children when compared to Prepubertal and Pubertal children. On the other hand, a significantly (P<0.05) lower reaction time as measured by ruler drop test (RDT) was noted in Postpubertal children when compared to Prepubertal and Pubertal children (Table 2).

**Table-2: Physical fitness variable of Prepubertal, Pubertal and Postpubertal children**

Variables	Prepubertal	Pubertal	Postpubertal
RDT (cm)	19.7 ± 2.3	16.8 ± 2.8*	15.3 ± 2.8*#
MSRT (cm)	10.9 ± 1.8	14.0 ± 1.3*	16.0 ± 1.6*#
SUT (in 30 sec)	15.5 ± 1.5	17.2 ± 1.2*	18.9 ± 1.3*#
SLJT (m)	1.5 ± 0.2	1.7 ± 0.3*	1.9 ± 0.3*#
GSTR (kg)	18.0 ± 3.6	24.7 ± 3.5*	31.5 ± 3.4*#
GSTL (kg)	17.0 ± 2.7	24.0 ± 3.3*	29.9 ± 3.1*#
30MAT (sec)	6.6 ± 0.3	5.7 ± 0.4*	5.1 ± 0.5*#
Peak power (watt)	496.9 ± 22.1	716.3 ± 20.6*	821.4 ± 21.3*#

All the values were expressed as mean and standard deviation (SD), n=50; ANOVA followed by multiple comparison tests; \*P<0.05 when compare to prepubertal age group, #P<0.05, when compare to pubertal age group; RDT = ruler drop test, MSRT= modified sit and reach test, SUT = sit ups test, SLJT= standing long jump test, GSTR= grip strength of right hand, GSTL= grip strength of left hand, 30MAT= 30 meter acceleration test.

Variables	Prepubertal	Pubertal	Postpubertal
RHR	64.3 ± 3.3	65.2 ± 3.6 <sup>NS</sup>	64.0 ± 2.9 <sup>NS</sup>
HRmax	195.3 ± 6.0	190.8 ± 5.7*	186.7 ± 6.4* <sup>#</sup>
RecHR1	158.6 ± 5.5	148.3 ± 6.1*	142.3 ± 4.7* <sup>#</sup>
RecHR2	134.6 ± 4.1	128.3 ± 3.5*	121.4 ± 4.4* <sup>#</sup>
RecHR3	115.1 ± 4.8	108.2 ± 2.5*	102.2 ± 2.7* <sup>#</sup>
RSBP	95.5 ± 5.0	99.8 ± 5.3*	102.9 ± 6.2* <sup>#</sup>
RDBP	60.2 ± 4.7	62.4 ± 4.9*	66.4 ± 4.1* <sup>#</sup>
VO <sub>2</sub> max (ml/kg/min)	37.8 ± 2.3	43.2 ± 3.1*	45.5 ± 3.7* <sup>#</sup>
PEFR (l)	224.0 ± 10.7	289.3 ± 12.4*	346.5 ± 12.1* <sup>#</sup>

All the values were expressed as mean and standard deviation (SD), n=50; ANOVA followed by multiple comparison tests; \*P<0.05 when compare to prepubertal age group, #P<0.05, when compare to pubertal age group, NS= not significant; RHR= resting heart rate, HRmax= maximal heart rate, RecHR1= recovery heart rate in 1<sup>st</sup> min, RecHR2= recovery heart rate in 2<sup>nd</sup> min, RecHR3= recovery heart rate in 3<sup>rd</sup> min, RSBP= resting systolic blood pressure, RDBP= resting diastolic blood pressure, VO<sub>2</sub>max= maximal aerobic capacity, PEFR= peak expiratory flow rate.

Cardiorespiratory fitness variables showed remarkable changes among the Prepubertal, Pubertal & Postpubertal children. A significantly (P<0.05) higher VO<sub>2</sub>max, peak expiratory flow rate (PEFR), resting systolic and diastolic blood pressure were observed in Postpubertal children when compared to Prepubertal and Pubertal children. On the other hand, a significantly (P<0.05) lower maximal heart rate (HRmax) and recovery heart rates were noted in Postpubertal children when compared to Prepubertal and Pubertal children. However, no significant change was reported in resting heart rate among the groups (Table 3).

### Discussion

Childhood and adolescence are crucial periods of life, since dramatic physiological and psychological changes take place at these ages. Physical growth in children is measured by changes in body size and/or composition as well as physical profile [4]. During childhood and adolescence, body size and composition markedly change. These changes are strongly associated with the development of various physical performance characteristics. At the same time, anthropometry and body composition during adolescence are predictors of risk factors for cardiovascular disease, diabetes, and many types of chronic diseases [17-18] which occur in adults [19-21]. Hence, determining anthropometry and body composition during childhood and adolescence would be of interest to those working in both sports sciences and medicine.

Body size (height, body mass, BMI and BSA) play important role during selection of players [22-24]. The tall players are recruited as in athletics, soccer, volleyball and other games. Although, game like field hockey has no significant impact on height; however a standard height should be maintained for selection of players for each sports discipline. Body mass is a considerable factor in games and sports, since body contact is essential in like soccer, field hockey and some other games [25-26]. Body mass index (BMI) has been used as a simple anthropometric index which reflects the current nutritional status of an individual, and that of body surface area (BSA) can be made of an individual's daily resting energy expenditure [4].

In the present study, a significantly higher height, body mass and BSA were observed in Postpubertal children when compared to Prepubertal and Pubertal children. However, no significant change was reported in BMI among the groups. The possible reason for the increase in height Prepubertal to Pubertal and Postpubertal children may be the osteotropic response to exercise. The osteotropic effect of exercise is dependent on load dynamics, the volume, intensity and duration of training, administered on the individual and the period in life when exposure occurs [27]. In addition, the hormonal regulation of skeleton is unique in each stage of life [28]. The gain in height is dependent on growth hormone and exercise is

a potent stimulus for growth hormone [28]. It has been reported that genetic influence can alter morphological status only within a narrow limit, set by his genotype [29].

Growth in body weight follows the same trend as in case of height. Increment in body weight in each age category may be due to the increment in bone and muscle weight. Increase in muscle mass with age, appears to result primarily from hypertrophy of existing fibres. The gain in weight is dependent on growth hormone and exercise is a potent stimulus for growth hormone [28]. Apart from the hormonal effects the neural maturity also helps to gain desirable body weight in the athletes [30]. It is possible that a particular body size will encourage acquisition of certain skills and force gravitation towards a specific playing position: this is likely to occur before maturity so that the individual will tend to favour one positional role before playing at senior level [25]. The waist to hip ratio (WHR) has been shown to be related to the risk of coronary heart disease [31-33]. Mid-upper arm circumference (MUAC) is a measure of nutritional status [31, 34].

The waist- hip ratio (WHR) of pubertal children was noted significantly higher ( $P<0.05$ ) when compared to prepubertal and postpubertal children. Moreover, a significantly higher mid upper arm circumference (MUAC) was observed in Postpubertal children when compared to Prepubertal and Pubertal children. The changes may be because of level of maturation factors and / or motivation, and exposure to long term and higher intensity of training among the Postpubertal children when compared with Prepubertal and Pubertal children. Similar findings were also noted by other research groups who reported significant change in these parameters with the advancement of age, level of maturation and exposure to high intensity of exercise for long time among the children [35-37]. Monitoring of anthropometry and body composition at regular intervals is essential for selection of athletes for competitions. In addition, the anthropometric variables can predict the risk of obesity, cardiovascular and other diseases.

The percentage of body fat plays an important role for the assessment of physical fitness of the players [25, 38-39]. Generally, the amount of fat in an adult male in his mid-twenties is about

16.5% of body weight [4, 30, 40]. A lean body is desirable for all sports discipline [25, 41-42]. A low-body fat may improve athletic performance by improving the strength-to-weight ratio [4, 30, 40]. Excess body fat adds to the load without contributing to the body's force-producing capacity [4, 30, 40]. A significantly lower ( $P<0.05$ ) percent body fat was observed in Postpubertal children when compared to Prepubertal and Pubertal children. The lower body fat values in the postpubertal children may be because of exposure to long term and higher intensity of aerobic endurance training compared with Prepubertal and Pubertal children. However, significant increase in LBM was noted in Postpubertal children when compared to Prepubertal and Pubertal children.

This may be again due to long term effect of exercise among the Postpubertal children than Prepubertal and Pubertal children which reduces the body fat and which shows higher LBM among the Postpubertal children [24, 40, 41]. It can be stated that excess body fat can limit the aerobic and anaerobic performance of the players. Similar observations have been noted by other research groups [24, 40, 41]. The observations of our study may be supported by several studies, where decrease in body fat was noted with the advancement of age of the players [24, 40]. Therefore, monitoring of body composition at regular intervals is essential for selection of athletes for competitions and during the training seasons.

Physical fitness of the athletes can be assessed by measuring motor skills and activities such as reaction time, hip and trunk flexibility, abdominal strength, elastic leg strength, grip strengths, maximum speed and peak power output. The reaction time is the time the athletes take for the body to react to a stimulus. The reaction time is very important for the track and field athletes as well as for players of different sports disciplines [1, 4]. In the present study, a significantly ( $P<0.05$ ) lower reaction time as measured by ruler drop test (RDT) was noted in Postpubertal children when compared to Prepubertal and Pubertal children. Flexibility is the ability to move a joint or series of joints smoothly and easily

throughout a full range of motion. An athlete who has a restricted range of motion will realize a decrease in performance capabilities. Flexibility is important in preventing injury to the musculotendinous and skeletal anatomy [1, 4]. There are some factors that limit flexibility are bony structure, excessive fat, skin, muscles and tendons, and connective tissues. With the exception of bony structure, age, and gender, all of the other factors that limit flexibility may be altered to increase range of joint motion. In the present study, a significantly ( $P < 0.05$ ) higher hip and trunk flexibility as measured by modified sit and reach test (MSRT) score was observed in Postpubertal children when compared to Prepubertal and Pubertal children. Strength is the central component of a athletics training program particularly for short distance run, through events, jump events and in different games [25, 39, 40].

Abdominal strength is important to monitor the development of the athlete's muscular endurance. To elastic leg strength is important to monitor the development of the athlete's muscular power. On the other hand, strength of grip muscle also has significant impacts on the performance of athletes, which is needed for throw-in, catching, serving, smashing or fisting the ball [25, 40]. In the present study, a significantly ( $P < 0.05$ ) higher abdominal strength as measured by sit ups test (SUT) score, elastic leg strength as measured by standing long jump test (SLJT) score and grip strengths of both hands were observed in Postpubertal children when compared to Prepubertal and Pubertal children. Assessment of speed is important for selection of athletes in teams. Speed relates to the ability to perform a movement within a short time period. Power is the amount of work done or energy transferred per unit of time. Muscular power is the ability to use strength quickly to produce an explosive effort. Sports like short distance run, soccer, field hockey etc. demands high speed and power output as quick acceleration and deceleration are important in this sport [25, 40].

Repeated back-to-back sprints make speed and tolerance to lactic acid an important characteristic in athletes [25, 40]. A high speed and power output are essential for such activities [25, 40]. Thus a high speed and power output helps to develop sprint quality of the athletes [25, 40]. In the present study, a significantly ( $P < 0.05$ ) higher

maximum speed as measured by lower 30 meter acceleration test (30MAT) score, and peak power output were observed in Postpubertal children when compared to Prepubertal and Pubertal children. The lower reaction time; and higher level of flexibility, abdominal strength, elastic leg strength and grip strength, speed and power of the postpubertal children may be because of exposure to long term and higher intensity of training when compared with Prepubertal and Pubertal children. Moreover, this may be because of level of maturation factors and / or motivation of the Postpubertal children when compared with Prepubertal and Pubertal children. Similar findings were also noted by other research groups who reported significant reduction of reaction time; and elevation in flexibility, abdominal strength, elastic leg strength and grip strength, speed and power with the advancement of age, level of maturation and exposure to high intensity of exercise for long time among the children [24, 35-37, 40]. Monitoring of the motor skills and activities such as reaction time, hip and trunk flexibility, abdominal strength, elastic leg strength, grip strengths, maximum speed and peak power output at regular intervals is essential for selection of athletes for competitions and during the training seasons.

Heart rate and blood pressure are essential for assessing cardiovascular fitness of the athletes. Heart rate increases with an increase in work intensity and shows a linear relationship with work rate [42]. The highest rate at which the heart can beat is the maximal heart rate (HRmax). Quick recovery from strenuous exercise is important in sports which involves intermittent efforts interspersed with short rests [4, 30, 43]. The heart rate recovery curve is an excellent tool for tracking a person's progress during a training program [4, 30]. A significantly ( $P < 0.05$ ) lower maximal heart rate (HRmax) and recovery heart rates were noted in Postpubertal children when compared to Prepubertal and Pubertal children. However, no significant change was reported in resting heart rate among the groups. On the other hand, a significantly ( $P < 0.05$ ) higher resting systolic and diastolic blood pressure were observed in Postpubertal children when

compared to Prepubertal and Pubertal children. Exercise cardio acceleration results from release of parasympathetic inhibition at low exercise intensities and from both parasympathetic inhibition and sympathetic activation at moderate intensities [4, 30]. Nevertheless, parasympathetic activation is considered to be the main mechanism underlying exponential cardio deceleration after exercise [4, 30]. The results of the present study suggest that the strain on the circulatory system during sports activities is relatively high. Exercising at this intensity should provide a good training stimulus. Therefore, heart rate and blood pressure monitoring is essential for selection of athletes and during the training seasons.

The maximal oxygen uptake ( $VO_{2max}$ ) is the best overall measure of aerobic power [37, 44]. Aerobic capacity certainly plays an important role in athletics activities and has a major influence on technical performance and tactical choices [4, 30, 40]. A significantly ( $P<0.05$ ) higher  $VO_{2max}$  was observed in Postpubertal children when compared to Prepubertal and Pubertal children. The higher level of  $VO_{2max}$  value in the postpubertal children may be because of exposure to long term and higher intensity of aerobic endurance training compared with Prepubertal and Pubertal children. The increase in  $VO_{2max}$  may be due to an increase in the systemic a-v  $O_2$  difference and stroke volume, [4, 30]. Moreover, these changes may be the result of increased volume of endurance training [4, 30]. The aerobic endurance training enhances the activity of the cardiovascular system as well as developed oxidative capacity of the skeletal muscles which leads to an increase in the delivery of oxygen to working muscles [4, 30]. This is accepted as the main reason for elevation of  $VO_{2max}$  (4, 30).

Similar observation has been reported previously [25, 40, 45]. The extent by which  $VO_{2max}$  could be changed with training also depends on the starting point [4, 30]. The fitter an individual is to begin with, the less potential there is for an increase and most elite athletes hit this peak early in their career [4, 30]. There also seems to be a genetic upper limit beyond which further increases in either intensity or volume have no effect on aerobic power [4, 30]. Other than tactical and technical aspects of soccer, monitoring of  $VO_{2max}$  is essential during the

training phases, which helps the coaches for selection of players for competition.

Lung function tests are of little value for predicting fitness and exercise performance, provided that the values fall within a normal range. Peak expiratory flow rate (PEFR) is used as an indicator of asthma or similar lung disease [4, 30]. In the present study, a significantly ( $P<0.05$ ) higher peak expiratory flow rate (PEFR) was observed in Postpubertal children when compared to Prepubertal and Pubertal children. The higher peak expiratory flow rate (PEFR) of the postpubertal children may be because of exposure to long term and higher intensity of aerobic endurance training when compared with Prepubertal and Pubertal children. Moreover, this may be because of level of maturation factors and / or motivation of the Postpubertal children when compared with Prepubertal and Pubertal children. Monitoring of PEFR at regular intervals is essential for selection of athletes for competitions and during the training seasons.

### Conclusion

Identification of children at early stage of their growth and development may produce elite athletes in the future. Talent identification also can be used as a counseling technique that helps to discover and explore areas of talent for particular athletes. In order to reach their goals, young children should be subject to a series of tests reflecting anthropometric, physical and cardiorespiratory fitness which will indicate their present over all strengths and weaknesses. Improvement in these parameters depends on level of maturation factors and / or motivation, and exposure to long term and higher intensity of training.

### Acknowledgment

The authors are indebted to the University Grant Commission, New Delhi, India for financial assistance to carry out the research work. The authors sincerely and wholeheartedly acknowledge the contribution of children participated in the present study. The authors are also thankful to the UG students, coaches and laboratory staffs for extending their support for the study.

## References

1. Reilly T, Secher N. Physiology of sports. *E & FN Spon: London*, 1990; 372-425.
2. Hoare DG. Predicting success in junior elite basket ball players-the contribution of anthropometric and physiological attributes. *J Sci Med Sports* 2000; 34: 391-405.
3. Coh M, Milanovic D, Embersic D. Anthropometric characteristics of elite junior male and female javelin throwers. *Coll Antropol.* 2002; 26:77-83.
4. Mc Ardle WD, Katch FI, Katch VL. Essentials of Exercise Physiology. 3<sup>rd</sup> ed. Lippincott Williams and Wilkins: *Philadelphia PA*, 2006.
5. Jonathan M, Mc Gavock BD, Torrance K, Mc Guire A, Wozny PD, Lewanczuk RZ. Cardiorespiratory Fitness and the Risk of Overweight in Youth: The Healthy Hearts Longitudinal Study of Cardiometabolic Health. *Obesity* 2009; 17:1802-7.
6. Ortega FB, Ruiz JR, Castillo MJ, Sjo"stro"m M. Physical fitness in childhood and adolescence: a powerful marker of health. *Int J Obesity* 2008; 32:1-11.
7. Payne N, Gledhill N, Katzmarzy KPT, Jamnik VK, Keir PJ. Canadian musculoskeletal norms. *Can J Appl Physiology* 2000; 25(6): 430-42.
8. Chatterjee S, Mandal A, Das NK. Physical and motor fitness level of Indian school going boys. *J Sports Med Phys Fitness* 1993; 33:268-77.
9. Manna I, Dhingra M, Khanna GL. Study of anthropometric and physiological profiles for the selection of under 14 years players in scheme. *J Sports Traumatol All Sports Sci* 2002; 4:53-8.
10. Manna I, Ghosh N, Banerjee S, Ghosh S, Kar SK, Dhara P. Effect of yoga on flexibility and reaction time in adolescent boys and girls. *Ind J Sports Studies* 2004; 3: 29-35.
11. Jonson BL, Nelson JK. Practical measurements for evaluation in physical education. *Macmillan Publishing Co: London*, 1996.
12. Durnin JVGA, Womersley J. Body fat assessed from total body density and its estimation from skin fold thickness: measurements on 481 men and women from 16 to 72 years. *Br J Nutr* 1974; 32:77-97.
13. Siri WE. The gross composition of the body. In: Tobias CA, Lawrence JH, ed. *Advances in Biological and Medical Physics. Academic Press: New York*, 1956; 239-80.
14. Margaria R, Aghemo P and Rovelli E. Measurement of muscular power (anaerobic) in man. *J Appl Physiol* 1996; 21:1662-64.
15. McArdle WD, Katch FI, Pechar GS, Jacobson L, Ruck S. Reliability and interrelationships between maximal oxygen intake, physical work capacity and step-test scores in college women. *Med Sci Sports* 1972; 4:182-86.
16. Mustajbegovic J, Zuskin E, Schachter EN, Kern J, Luburic-Milas M, Pucarin J. Respiratory findings in tobacco workers. *Chest* 2003; 123:1740-48.
17. Dietz WH. Childhood weight affects adult morbidity and mortality. *J Nutr* 1998; 128:411S-14S.
18. Goran MI, Ball GD, Cruz ML. Obesity and risk of type 2 diabetes and cardiovascular disease in children and adolescents. *J Clin Endocrinol Metab* 2003;88:1417-27.
19. Guo SS, Chumlea WC, Roche AF, Siervogel RM. Age-and maturity-related changes in body composition during adolescence into adulthood: the Fels longitudinal study. *Int J Obesity* 1997; 21:1167-75.
20. Katzmarzyk PT, Perusse L, Malina RM, Bergeron J, Despres JP, Bouchard C. Stability of indicators of the metabolic syndrome from childhood and adolescence to young adulthood: the Quebec family study. *J Clin Epidemiol* 2001; 54:190-95.
21. Janssen I, Katzmarzyk PT, Srinivasan SR, Chen W, Malina RM, Bouchard C, Berenson GS. Utility of childhood BMI in the prediction of adulthood disease: comparison of national and international references. *Obes Res* 2005; 13:1106-15.
22. Enemark-Miller EA, Seegmiller JG, Rana SR. Physiological profile of women's Lacrosse players. *J Strength Cond Res* 2009; 23:39-43.
23. Johnson A, Doherty PJ, Freemont A. Investigation of growth, development, and factors associated with injury in elite schoolboy footballers: prospective study. *BMJ* 2009; 338: b490.
24. Silvestre R, West C, Maresh CM, et al. Body composition and physical performance in men's soccer: a study of a National Collegiate Athletic Association Division I team. *J Strength Cond Res* 2006; 20:177-83.
25. Hoff J. Training and testing physical capacities for elite soccer players. *J Sports Sci* 2005; 23:573-82.
26. Tahara Y, Moji K, Tsunawake N et al. Physique, body composition and maximum oxygen consumption of selected soccer players of Kunimi High School, Nagasaki, Japan. *J Physiol Anthropol* 2006; 25:291-7.
27. Mujika I. The influence of training characteristics and tapering on the adaptation in highly trained individuals: a review. *Int J Sports Med* 1998; 19:439-46.
28. Coyel EF, Hemmert MK, Coggan AR, et al. Effect of detraining on cardiovascular responses to exercise: role of blood volume. *J Appl Physiol* 1986; 60:95-9.
29. Reilly T, Williams AM, Nevil A, Franks A. A multidisciplinary approach to talent identification in soccer. *J Sports Sci* 2000; 18:695-702.
30. Wilmore JH, Costill DL. Physiology of Sport and Exercise. 3<sup>rd</sup> ed. *Human Kinetics: Champaign IL*, 2005.
31. Bassareo PP, Marras AR, Barbanti C, Mercurio G. Comparison between waist and mid-upper arm circumferences in influencing systolic blood pressure in adolescence: the SHARP (Sardinian Hypertensive Adolescent Research Programme) study. *J Pediatr Neonat Individual Med* 2013; 2(2): e020207.
32. Dobbeltsteyn CJ et al. A comparative evaluation of waist circumference, waist-to-hip ratio and body mass index as indicators of cardiovascular risk factors. The Canadian Heart Health Surveys. *Int J Obes Relat Metab Disord* 2001; 25(5):652-61.



33. Yalcin BM, Sahin EM, Yalcin E. Which anthropometric measurement is most closely related to elevated blood pressure? *Fam Pract* 2005; 22:541-47.
34. James WPT, Mascie-Taylor CGN, Norgan NG, Bristrian BR, Shetty P, Ferro-Luzzi A. The value of arm circumference measurements in assessing chronic energy deficiency in Third World adults. *Eur J Clin Nutr* 1994; 48:883-94.
35. Fukunaga Y, Takai Y, Yoshimoto T, Fujita E, Yamamoto M, Kanehisa H. Influence of maturation on anthropometry and body composition in Japanese junior high school students. *J Physiol Anthropol* 2013; 32(1):5.
36. Ujević T, Sporis G, Milanović Z, Pantelić S, Neljak B. Differences between health-related physical fitness profiles of Croatian children in urban and rural areas. *Coll Antropol* 2013; 37(1):75-80.
37. Coelho-E-Silva MJ, Vaz Ronque ER, Cyrino ES, et al. Nutritional status, biological maturation and cardiorespiratory fitness in Azorean youth aged 11-15 years. *BMC Public Health* 2013; 13: 495.
38. Ostojic SM. Seasonal alterations in body composition and sprint performance of elite soccer players. *J Exer Physiol on line* 2003; 6:24-7.
39. Svensson M, Drust B. Testing soccer players. *J Sports Sci* 2005; 23:601-18.
40. Reilly T. An Ergonomics model of the soccer training process. *J Sports Sci* 2005; 6:561-72.
41. Ekelund U, Poorvliet E, Nilson A, et al. Physical activity in relation to aerobic fitness and body fat in 14- to 15- year-old boys and girls. *Eur J Appl Physiol* 2001; 85:195-201.
42. Astrand PO, Rodhal K. Textbook of work physiology. *McGraw-Hill: New York*, 1986.
43. Rampinini E, Impellizzeri FM, Castagna C et al. Factors influencing physiological responses to small-sided soccer games. *J Sports Sci* 2007; 25:659-66.
44. Popadic Gacesa JZ, Barak OF, Grujic NG. Maximal anaerobic power test in athletes of different sport disciplines. *J Strength Cond Res* 2009; 23:751-55.
45. Miller TA, Thierry-Aguilera R, Congleton JJ, et al. Seasonal changes in VO<sub>2</sub>max among Division 1A collegiate women soccer players. *J Strength Cond Res* 2007; 21:48-51.

\*All correspondences to: Dr. Indranil Manna, Assistant Professor, Department of Physiology, Midnapore College, Raja Bazar Main Road, Midnapore-721101, West Bengal, India. E-mail ID: indranil\_manna@yahoo.com